

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/16, 38/18, 38/30, 38/17, 38/39 // (A61K 38/16, 39:395, 38:17, 31:70)		A1	(11) International Publication Number: WO 99/55359
			(43) International Publication Date: 4 November 1999 (04.11.99)
(21) International Application Number: PCT/US99/08303 (22) International Filing Date: 15 April 1999 (15.04.99) (30) Priority Data: 60/083,472 29 April 1998 (29.04.98) US (71) Applicant: ALLERGAN SALES, INC. [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US). (72) Inventors: DOLLY, J., Oliver; 7 Buckland Road, Cheam, Surrey SM2 7LP (GB). AOKI, Kei, Roger; 2 Ginger Lily Court, Coto de Caza, CA 92679 (US). DE PAIVA, Anton; 62 Ravensmede Way, London W4 1TF (GB). (74) Agents: FISHER, Carlos, A. et al.; Allergan Sales, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: COMPOSITIONS AND METHODS FOR EXTENDING THE ACTION OF CLOSTRIDIAL NEUROTOXIN			
(57) Abstract Methods and compositions for modulating neurite outgrowth in damaged neural endplates. Also disclosed are methods for extending the period during which tissue treated with Clostridial toxin is paralysed.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOSITIONS AND METHODS FOR EXTENDING THE ACTION OF CLOSTRIDIAL NEUROTOXIN

This application claims priority under 35 U.S.C.
5 §119(e)(1) to provisional application number 60/083,472,
filed 4/29/98.

Field of the Invention

10 The present invention is directed towards methods and
compositions for inhibiting neural sprouting in neurons that
have been subjected to botulinum toxin. Also disclosed are
methods and compositions for extending the period of time
during which treatment of nerve cells with botulinum toxin
15 is effective to prevent innervation of a cell or tissue,
such as muscle cells or tissue. Such methods and
compositions are effective in the treatment of spasms or
muscular tetanus. Disclosed as well are methods and
compositions for stimulating neural outgrowth.

20

Background of the invention

Neurotoxins, such as those obtained from *Clostridium*
botulinum and *Clostridium tetanus*, are highly potent and
specific poisons of neural cells. These Gram positive
25 bacteria secrete two related but distinct toxins, each
comprising two disulfide-linked amino acid chains: a light
chain (L) of about 50 KDa and a heavy chain (H) of about 100
KDa, which are wholly responsible for the symptoms of these
diseases.

The tetanus and botulinum toxins are among the most lethal substances known to man, having a lethal dose in humans of between 0.1 ng and 1 ng per kilogram of body weight. Tonello et al., *Adv. Exp. Med. & Biol.* 389:251-260 (1996). Both toxins function by inhibiting neurotransmitter release in affected neurons. The tetanus neurotoxin (TeNT) acts mainly in the central nervous system, while botulinum neurotoxin (BoNT) acts at the neuromuscular junction by inhibiting acetylcholine release from the axon of the affected neuron into the synapse, resulting in a localized flaccid paralysis. The effect of intoxication on the affected neuron is long-lasting and has been thought to be irreversible.

The tetanus neurotoxin (TeNT) is known to exist in one immunologically distinct type; the botulinum neurotoxins (BoNT) are known to occur in seven different immunogenic types, termed BoNT/A through BoNT/G. While all of these types are produced by isolates of *C. botulinum*, two other species, *C. baratii* and *C. butyricum* also produce toxins similar to /F and /E, respectively. See e.g., Coffield et al., *The Site and Mechanism of Action of Botulinum Neurotoxin in Therapy with Botulinum Toxin* 3-13 (Jankovic J. & Hallett M. eds. 1994), the disclosure of which is incorporated herein by reference.

Regardless of type, the molecular mechanism of intoxication appears to be similar. In the first step of the process, the toxin binds to the presynaptic membrane of the target neuron through a specific interaction between the

heavy chain and a cell surface receptor; the receptor is thought to be different for each type of botulinum toxin and for TeNT. The carboxy terminus of the heavy chain appears to be important for targeting of the toxin to the cell surface.

In the second step, the toxin crosses the plasma membrane of the poisoned cell. The toxin is first engulfed by the cell through receptor-mediated endocytosis, and an endosome containing the toxin is formed. The toxin then escapes the endosome into the cytoplasm of the cell. This last step is thought to be mediated by the amino terminus of the heavy chain, which triggers a conformational change of the toxin in response to a pH of about 5.5 or lower. Endosomes are known to possess a proton pump which decreases intra-endosomal pH. The conformational shift exposes hydrophobic residues in the toxin, which permits the toxin to embed itself in the endosomal membrane. The toxin then translocates through the endosomal membrane into the cytosol.

The last step of the mechanism of botulinum toxin activity appears to involve reduction of the disulfide bond joining the heavy and light chain. The entire toxic activity of botulinum and tetanus toxins is contained in the light chain of the holotoxin; the light chain is a zinc (Zn++) endopeptidase which selectively cleaves proteins essential for recognition and docking of neurotransmitter-containing vesicles with the cytoplasmic surface of the plasma membrane, and fusion of the vesicles with the plasma

membrane. TxNT, BoNT/B, BoNT/D, BoNT/F, and BoNT/G cause degradation of synaptobrevin 2 (also called vesicle-associated membrane protein (VAMP)), a synaptosomal membrane protein. Most of the VAMP present at the cytosolic surface of the synaptic vesicle is removed as a result of any one of these cleavage events. Each toxin specifically cleaves a different bond.

BoNT/A and /E selectively cleave the plasma membrane-associated protein SNAP-25; this protein is bound to and present on the cytosolic surface of the plasma membrane. BoNT/C cleaves syntaxin, an integral protein having most of its mass exposed to the cytosol. Syntaxin interacts with the calcium channels at presynaptic terminal active zones. See Tonello et al., *Tetanus and Botulism Neurotoxins in Intracellular Protein Catabolism* 251-260 (Suzuki K & Bond J. eds. 1996), the disclosure of which is incorporated by reference as part of this specification. BoNT/C₁ also cleaves SNAP-25 at a peptide bond next to that cleaved by BoNT/A.

Both TeNT and BoNT are taken up at the neuromuscular junction. BoNT remains within peripheral neurons, and blocks release of the neurotransmitter acetylcholine from these cells. Through its receptor, TeNT enters vesicles that move in a retrograde manner along the axon to the soma, and is discharged into the intersynaptic space between motor neurons and the inhibitory neurons of the spinal cord. At this point, TeNT binds receptors of the inhibitory neurons, is again internalized, and the light chain enters the

cytosol to block the release of the inhibitory neurotransmitters 4-aminobutyric acid (GABA) and glycine from these cells. Id.

Because of its specifically localized effects, dilute
5 preparations of BoNT have been used since 1981 as
therapeutic agents in the treatment of patients having
various spastic conditions, including strabismus
(misalignment of the eye), buphthalmos (involuntary eyelid
closure) and hemifacial spasm. See e.g., Borrodi et al.,
10 *Pharmacology and Histology Botulinum Toxin in Therapy with
Botulinum Toxin* 3-13 (Jankovic J. & Hallett M. eds. 1994),
incorporated by reference herein. Of the seven toxin types,
BoNT/A is the most potent of the BoNTs, and the most well
characterized. Intramuscular injection of spastic tissue
15 with dilute preparations of BoNT/A has been also used
effectively to treat spasticity due to brain injury, spinal
cord injury, stroke, multiple sclerosis and cerebral palsy.
The extent of paralysis depends on both the dose and dose
volume delivered to the target site. Typically, the
20 neurotoxin is administered in a preparation that also
contains several non-toxic proteins as well, including
hemagglutinins and associated glycoproteins that assist in
maximizing its stability and presentation to the target
motor neuron.

25 Typically there is a 24 to 72 hour delay between the
administration of the toxin and the onset of the clinical
effect. Exposure to the toxin causes denervation atrophy.
See e.g., Dutton J., *Acute and Chronic Effects of Botulinum*

Toxin in the Management of Blepharospasm, in Therapy with Botulinum Toxin at 199, incorporated by reference herein.

Although the therapeutic application of BoNT has been extraordinarily effective, what side effects have been
5 observed are mainly immediate effects associated with the treatment event itself. Peripheral effects causing weakness in adjacent muscles may sometimes occur; these effects usually do not persist beyond 1 to two weeks. The specific manifestations of these effects upon adjacent muscle groups
10 depend upon the particular indication being treated. For example, patients treated for blepharospasm sometimes experience ptosis, and swallowing problems may occur after injection of neck muscles for torticollis.

Other possible consequences of treatment include the
15 potential for overdosage through miscalculation or differences in activity between different preparations of the toxin, generalized fatigue, and the potential for allergic reactions.

A feature of treatment with BoNT/A, and other
20 clostridial neurotoxin types, is that the paralytic action is temporary with symptoms reappearing in patients within a few months after toxin injection. This characteristic has been thought to be associated with the observed sprouting of nascent, synaptically active processes at the neuromuscular
25 junction (NMJ). The production of such sprouts following BoNT/A therapeutic treatment has appeared to contribute to the reinnervation of the treated tissue and therefore the need for repeated serial injections of the toxin.

The duration of paralytic action caused by clostridial neurotoxin and the extent of sprouting, appears to depend on the neurotoxin subtype studied and therefore appears to be related to the specific SNARE target cleaved by each neurotoxin. Thus, BoNT/A and BoNT/C₁, which cleave the t-SNARE protein SNAP-25 within one amino acid of each other, cause a long lasting paralysis with a long average sprout length observed. BoNT/F, which cleaves the v-SNARE protein VAMP, causes a paralysis of shorter duration and sprouts of shorter average length. BoNT/E, which cleaves SNAP-25 at a different position than that of BoNT/A and BoNT/C₁ (thereby liberating a SNAP fragment of different size from the plasma membrane), has a short duration period of about 5 days, and virtually no neural sprouting is observed. Without wishing to be bound by theory, these observations suggest that neural sprouting and duration of paralysis are normally related events, and that the cleavage products of BoNT proteolytic digestion (i.e., the liberated fragment or the membrane-bound fragment) can directly or indirectly regulate, or be coregulated with, neural sprouting.

It would therefore be advantageous to design a method whereby the sprouting phenomenon may be uncoupled from duration of therapeutic effect, and thus delayed, blocked or attenuated so as to prolong the effects of injection of tissue with toxin. Although the experiments described herein utilize a BoNT/A preparation, it will be recognized by those of skill in the art that the methods shown herein will be suitable for employment using other clostridial

toxins, such as BoNT/B through /G, and TeNT, in which a sprouting pathway can be observed.

Additionally, it would be advantageous to provide herein compositions effective for the inhibition or
5 prevention of the sprouting phenomenon. Such compositions and methods would lessen the need for patients to undergo repeated neurotoxin treatment.

Summary of the Invention

10

The present invention concerns methods for increasing the period of time between therapeutic treatments of neural tissue with a clostridial neurotoxin; thus the method provides a method of increasing the effectiveness of such
15 treatments. A direct advantage of such methods is an increased therapeutic "life" and a concomitant lessening in the required frequency of treatment of the patient with neurotoxin. Reducing frequency of treatment would provide less opportunity for a patient to experience the side
20 effects described above that are observed following treatment, but which tend to subside long before the effectiveness of the toxin in the target area has subsided. Additionally, reduced frequency of treatment provides less opportunity for miscalculation of dosage amount and other
25 treatment-specific risks.

Accordingly, an aspect of the invention concerns a composition comprising a first agent comprising a clostridial neurotoxin for use as a therapeutic agent and a

second agent able to extend the duration of therapeutic benefit of said first agent, wherein the second agent is effective to attenuate the production of nerve terminal sprouts following treatment of a neuromuscular junction with
5 the clostridial neurotoxin.

In a particular embodiment of this aspect of the invention, the first and second agent may comprise a single entity which is provided the patient in a single treatment session. For example, the entity may comprise a single
10 molecule, or a disulfide-linked multichain polypeptide. Additionally or alternatively, the entity may comprise one or more adsorbed or linked heterogroup, such as a small organic molecule or a nucleic acid linked thereto. The entity preferably comprises both the receptor binding and
15 translocation activities of a clostridial heavy chain and an active portion of a clostridial toxin light chain. The light chain may also comprise an auxiliary enzymatic activity, such as a ribonuclease, which specifically cleaves a nucleic acid encoding an intraneuronal factor which is
20 responsible for the expression, activation and/or secretion of neurotrophic factors or cell adhesion molecules. In a preferred embodiment, such an auxiliary activity is provided by a ribozyme. By ribozyme is meant a nucleic acid or nucleic acid analog having a sequence-specific nuclease
25 activity; the construction and use of ribozymes are well known in the art; see e.g., Cech, T., *Science* 236:1532-1539(1987); Cech, T. R., *Curr. Opin. Struct. Biol.* 2:605-609 (1992); and Usman et al., *Nucleic Acids & Mol. Biol.* 10:243-

264 (1996), the disclosures of which are hereby incorporated by reference herein. By nucleic acid analog is meant a polymeric molecule able to form a sequence-specific hybrid with a target single-stranded nucleic acid; such analogs may
5 contain modified nucleotides (or ribonucleotides) such as 3'-O methyl nucleotides, phosphorothioate modified nucleotides, methylphosphonate nucleotides, or nucleotide bases separated by a peptide-like bond.

Alternatively, a nucleic acid or nucleic acid analog
10 comprised in the single entity referred to above may be an antisense agent able to selectively bind to a nucleic acid encoding an intraneuronal factor which is responsible for the expression, activation and/or secretion of neurotrophic factors or cell adhesion molecules. This antisense agent
15 may further provide a double-stranded substrate for the action of an intracellular RNase H activity. Details concerning certain embodiments of these aspects of the invention are contained in e.g., Dolly et al., International Publication No. WO95/32738, entitled *Modification of*
20 *Clostridial Toxins for Use as Transport Proteins* and Uherek et al., *J. Biol. Chem.* 273:8835-8841 (1998). These two references are incorporated by reference as part of the present application.

A nucleic acid moiety linked polypeptide portion of the
25 entity may encode a protein or polypeptide having the ability to be expressed within a neuron and to directly or indirectly regulate the expression, activation and/or secretion of neurotrophic factors or cell adhesion

molecules.

In preferred aspects of the invention, the second agent is selected from the group consisting of agents able to compete with, down-regulate, or neutralize the effects of:

5 IGFI, IGF II, a neurotrophic factor, leukemia inhibitory factor, a nerve cell adhesion molecule and neural agrin. In a more preferred aspect of the invention, the neurotrophic factor is selected from the group consisting of: ciliary neurotrophic factor, NT-3, NT-4, and brain-derived

10 neurotrophic factor and/or the nerve cell adhesion molecule is selected from the group consisting of tenascin-C, ninjurin, neural cell adhesion molecule.

Applicants have surprisingly discovered that recovery of neural function following poisoning of nerve terminals

15 with clostridial neurotoxin involves two distinct and apparently coordinated events. First, the poisoned endplate becomes synaptically inactive. Shortly thereafter the endplate elaborates thin nascent axon neural processes. These processes or "sprouts" are synaptically competent

20 after about 14 days following treatment with clostridial neurotoxin. The sprouts continue growing, reaching a maximal length and level of complexity after about 42 days following treatment with neurotoxin. During this time, the endplate remains synaptically inactive.

25 Secondly, after about 42 days, the sprouts begin to regress, shortening in length and decreasing in complexity.

At the same time, the original endplate begins to become synaptically active, undergoing synaptic vesicle turnover.

The increase in such turnover reaches that of the unpoisoned endplate after about 91 days post-treatment, at approximately the same time that the sprouting phenomenon has completely regressed, and no sprouts can be observed.

- 5 These findings are reported in DePaiva et al., *Proc. Nat'l. Acad. Sci.* 96:3200-3205 (March 1999), which is hereby incorporated by reference herein.

- These observations are diametrically opposed to the prevailing wisdom in the art, in which it has largely been
10 assumed that the original endplate is permanently inactivated upon treatment with clostridial toxin, and that all return of synaptic activity is due to extension of sprouts to compensate for the lack of a neurologically functional endplate. However, as indicated, the old
15 paradigm has been shown by the present Applicants to be in error, in that the poisoned endplate regains neurological function over time, while the axon sprouts regress so that after a given time period the nerve terminal appears essentially as it did prior to treatment. Therefore,
20 Applicants have discovered that, far from being a permanent feature, the axonal sprouts assume a temporary role in the rehabilitation of the poisoned endplate.

- Although not wishing to be limited by theory, Applicants believe that these results indicate that the two
25 events outlined above are temporally coordinated, in that a blockage of the neural sprouting phenomenon would delay or block the recovery of the functional endplate. Such temporal coordination could be due to the secretion of one

or more factor by the damaged neural endplate (or the inactive muscle fiber) that has neurotrophic effects resulting in the formation of neural sprouts; these sprouts may then elaborate a factor (either the same or different
5 from the first factor) which promotes continued sprouting. This factor may be produced during neurite sprouting in amounts sufficient for the reinnervation of the neurotoxin-damaged endplate even after treatment with clostridial toxin. Thus, treatment of cells with an agent able to block
10 neural sprouting would also delay or otherwise attenuate the ability of the treated endplate to experience return of neurological function, and in fact may well block such return altogether.

As indicated, the signaling event indicating the
15 initiation of the neural sprouting phenomenon appears to be mediated by a cytokine or other intercellular messenger. One such agent, agrin, appear to be an important player, if not the key molecule, in the formation of the neuromuscular junction in development, and in neuromuscular regeneration.
20 See Ruegg M.A. and Bixby J. L., *Trends in Neurol. Sci.* 21:22-27 (1998), the disclosure of which is incorporated herein by reference. Agrin appears to be present in a number of isoforms, which result from alternative mRNA splicing. Soluble agrin isolated from synaptic basal lamina extracts
25 (to which it binds following secretion) is able to induce the aggregation of acetylcholine receptors in the postsynaptic portions of muscle cells. Agrin present in motor neuron terminals (n-agrin) contains an insert,

relative to other agrin species, in a region termed the B/z-region; this insert is important in conferring the ability on n-agrin to aggregate acetylcholine receptors in postsynaptic tissue. Neural agrin is released by the motor-
5 nerve terminal and is believed by Applicants to induce postsynaptic specialization and up-regulation of other factors, such as muscle-diffusable factors, involved in the neural sprouting response.

It is anticipated that methods which interfere with the
10 sprouting phenomenon (such as by the preventing the action of agrin) would delay the return of innervation to cells which are controlled by neurons which have been therapeutically poisoned (i.e., by BoNT or TeNT) or otherwise damaged. This is because, as indicated above, the
15 rate of return of neural function appears to be partly dependent upon the presence of synaptically active sprouts. Additionally, agents to be used in the inhibition of sprouting may very well also delay or prevent the recovery of neural activity of the endplate following poisoning.

20 Thus utilization of such methods would therefore be expected to extend the effective period of treatment of tissue with a clostridial toxin, by delaying the regeneration of active neuromuscular synapses, both through inhibition of sprouting and of recovery of the poisoned
25 endplate.

Detailed Description of the Invention

This invention is drawn to methods and compositions for increasing the therapeutic effectiveness of treatment of tissue with clostridial neurotoxin. This increase in effectiveness is made possible by the surprising discovery
5 that regeneration of neural tissue damaged by treatment with clostridial neurotoxin is a complex occurrence in which two coordinated events take place.

In the first of these events, the poisoned neuromuscular endplate becomes synaptically inactive,
10 demonstrating no exocytosis of synaptic vesicles and thus no transport of intracellular acetylcholine. In four days after treatment, the endplate begins to form neural sprouts that are shown to release and regenerate synaptic vesicles.

These sprouts grow in length and complexity until
15 approximately 42 days following treatment with the neurotoxin; at this point the neural sprouts begin to regress and shorten. At ninety-one days following treatment, the neural sprouts can no longer be seen.

In the second event, simultaneously with the beginning
20 of regression of the neural sprouts, the synaptically inactive endplate begins to regain the ability to release acetylcholine and begin to recycle synaptic vesicles. This ability, which begins at relatively low levels, increases over the time period indicated above. At approximately 91
25 days following treatment with clostridial neurotoxin the endplate is histologically and synaptically indistinguishable from the condition of the endplate before treatment with clostridial neurotoxin.

These findings indicate that one may therapeutically intervene at one of the major steps of the sprouting phenomenon to prevent or attenuate the neural sprouting as a method of extending the effective period during which tissue
5 treated with the toxin remains paralyzed. In an initial step, the muscle cells surrounding the neural endplate respond, either sensing the inactive muscle or in response to a signal from the poisoned nerve terminal, by producing muscle-derived diffusable factors. The expression of a
10 number of muscle-derived signaling factors appears to be upregulated by muscle inactivation; such factors include insulin-like growth factors (IGF-1 and IGF-2). A factor such as neural agrin is believed to be the initial signal directing the muscle cell to produce the IGF molecules.
15 Reports have demonstrated that IGF I and IGF II effect neurite outgrowth in cultured BoNT/A treated dorsal root ganglia, and also are able to stimulate the initial sprouting response in paralyzed mouse gluteus muscle. See Caroni, P. and Schneider, C. *J. Neurosci.* 14:3378-3388
20 (1994) and Caroni, P., et al. *J. Cell Biol.* 125:893-902 (1994).

Thus, blocking the effects of such muscle derived diffusable factors that positively affect neurite outgrowth and sprouting may attenuate not only clostridial neurotoxin-
25 induced sprouting, but may also delay the eventual recovery of neurotransmission at the poisoned nerve terminals. Such blocking may occur through the use of antibodies specific for the muscle derived diffusable factor in question, or

that are common to such muscle derived diffusable factors. .
Alternatively, there are naturally occurring binding
proteins, such as the IGF binding proteins IGF-BP 4 and IGF-
BP 5, which can bind to, and therefore block, the
5 neurotrophic effect of such diffusable factors.

IGF-BP 4 has an amino acid sequence (from the amino
terminus) of:

MLPLCLVAALLLAAGPGPSLGDEAIHCPPCSEEKLARCRPPVGCEELVREPGCGCCATCA
10 LGLGMPGCVYTTPRCGSLRCYPPRGVEKPLHTLMHGQGVCMELAEIEAIQESLQPSDKDE
GDHPNNSFSPCSAHDRLCLQKHFAKIRDRSTSGGKMKVNGAPREDARVPVQSGSCQSELHR
ALERLAASQSRTHEDLYIIPNCDRNGNFHPKQCHPALDGQRGKCWCVDRKTGVKLPGG
LEPKGELDCHQLADSFRE

15 (SEQ ID NO: 1)

This is encoded within the nucleotide base sequence (from 5'
to 3') of:

20 GTGCCCTCCG CCGCTCGCCC GCGCGCCCGC GCTCCCCGCC TCGCCCCAGC
GCCCCGCGCC CGCGCCCCAG TCCTCGGGCG GTCATGCTGC CCCTCTGCCT
CGTGGCCGCC CTGCTGCTGG CCGCCGGGCC CGGGCCGAGC CTGGGCGACG
AAGCCATCCA CTGCCC GCCC TGCTCCGAGG AGAAGCTGGC GCGCTGCCCG
CCCCCGTGG GCTGCGAGGA GCTGGTGCGA GAGCCGGGCT GCGGCTGTTG
25 CGCCACTTGC GCCCTGGGCT TGGGGATGCC CTGCGGGGTG TACACCCCCC
GTTGCGGCTC GGGCCTGCGC TGCTACCCGC CCCGAGGGGT GGAGAAGCCC
CTGCACACAC TGATGCACGG GCAAGGCGTG TGCATGGAGC TGGCGGAGAT
CGAGGCCATC CAGGAAAGCC TGCAGCCCTC TGACAAGGAC GAGGGTGACC
ACCCCAACAA CAGCTTCAGC CCCTGTAGCG CCCATGACCG CAGGTGCCTG
30 CAGAAGCACT TCGCCAAAAT TCGAGACCGG AGCACCAGTG GGGGCAAGAT
GAAGGTCAAT GGGGCGCCCC GGGAGGATGC CCGGCCTGTG CCCCAGGGCT
CCTGCCAGAG CGAGCTGCAC CGGGCGCTGG AGCGGCTGGC CGCTTCACAG
AGCCGCACCC ACGAGGACCT CTACATCATC CCCATCCCCA ACTGCGACCG
CAACGGCAAC TTCCACCCCA AGCAGTGTCA CCCAGCTCTG GATGGGCAGC
35 GTGGCAAGTG CTGGTGTGTG GACCGGAAGA CGGGGGTGAA GCTTCCGGGG
GGCCTGGAGC CAAAGGGGGA GCTGGACTGC CACCAGCTGG CTGACAGCTT
TCGAGAGTGA GGCCTGCCAG CAGGCCAGGG ACTCAGCGTC CCCTGCTACT
CCTGTGCTCT GGAGGCTGCA GAGCTGACCC AGAGTGGAGT CTGAGTCTGA

```

GTCCTGTCTC TGCCTGCGGC CCAGAAGTTT CCCTCAAATG CGCGTGTGCA
CGTGTGCGTG TGCCTGCGTG TGTGTGTGTT TGTGAGCATG GGTGTGCCCT
TGGGGTAAGC CAGAGCCTGG GGTGTTCTCT TTGGTGTTAC ACAGCCCAAG
AGGACTGAGA CTGGCACTTA GCCCAAGAGG TCTGAGCCCT GGTGTGTTTC
5 CAGATCGATC CTGGATTAC TCACTCACTC ATTCCTTAC TCATCCAGCC
ACCTAAAAAC ATTTACTGAC CATGTACTAC GTGCCAGCTC TAGTTTTCAG
CCTTGGGAGG TTTTATTCTG ACTTCCTCTG ATTTTGGCAT GTGGAGACAC
TCCTATAAGG AGAGTTCAAG CCTGTGGGAG TAGAAAAATC TCATTCCCAG
AGTCAGAGGA GAAGAGACAT GTACCTTGAC CATCGTCCTT CCTCTCAAGC
10 TAGCCAGAGG GTGGGAGCCT AAGGAAGCGT GGGGTAGCAG ATGGAGTAAT
GGTCACGAGG TCCAGACCCA CTCCCAAAGC TCAGACTTGC CAGGCTCCCT
TTCTCTTCTT CCCCAGGTCC TTCTTTAGG TCTGGTTGTT GCACCCTGTG
CTTGGTTGGC TGGCAGCTGA GAGCCCTGCT GTGGGAGAGC GAAGGGGGTC
AAAGGAAGAC TTGAAGCACA GAGGGCTAGG GAGGTGGGGT ACATTTCTCT
15 GAGCAGTCAG GGTGGGAAGA AAGAAATGCAA GAGTGGACTG AATGTGCCTA
ATGGAGAAGA CCCACGTGCT AGGGGATGAG GGGCTTCCTG GGTCTGTTC
CCTACCCCAT TTGTGGTCAC AGCCATGAAG TCACCGGGAT GAACCTATCC
TTCCAGTGGC TCGCTCCCTG TAGCTCTGCC TCCCTCTCCA TATCTCCTTC
CCCTACACCT CCCTCCCCAC ACCTCCCTAC TCCCCTGGGC ATCTTCTGGC
20 TTGACTGGAT GGAAGGAGAC TTAGGAACCT ACCAGTTGGC CATGATGTCT TTTCT

```

(SEQ ID NO: 3)

IGFBP 5 has an amino acid sequence (from the amino
 25 terminus) of:

```

MVLTLAVLLLLLAAYAGPAQSLGSFVHCEPCDEKALSMCPPSPLGCELVKEPGCGCCMTCA
LAEGQSCGVYTERCAQGLRCLPRQDEEKPLHALLHGRGVCLNEKSYREQVKIERDSREHE
EPTTSEMAEETYSPIFRPKHTRISELKAEAVKKDRRKLTQSKFVGGAENTAHPRIISA
30 PEMRQESEQGPCRRHMEASLQELKASPRMVPRAVYLPNCDRKGFKRKQCKPSRGRKRG
CWCVDKYGMKLPMEYVDGDFQCHTFDSSNVE

```

(SEQ ID No. 2)

35 and is encoded within a nucleotide base sequence (from 5' to
 3') of:

```

GGGGAAAAGA GCTAGGAAAG AGCTGCAAAG CAGTGTGGGC TTTTCCCTT
TTTTTGCTCCT TTTCATTAC CCCTCCTCCG TTTTCACCCT TCTCCGACT
40 TCGCGTAGAA CCTGCGAATT TCGAAGAGGA GGTGGCAAAG TGGGAGAAAA

```

```

GAGGTGTTAG GGTTTGGGGT TTTTTGTTT TTGTTTTTGT TTTTAAATTT
CTTGATTTCA ACATTTTCTC CCACCCTCTC GGCTGCAGCC AACGCCTCTT
ACCTGTTCTG CGGCGCCGCG CACCGCTGGC AGCTGAGGGT TAGAAAGCGG
GGTGTATTTT AGATTTTAAG CAAAAATTTT AAAGATAAAT CCATTTTTCT
5 CTCCACCCCC CAACGCCATC TCCACTGCAT CCGATCTCAT TATTCGGTG
GTTGCTTGGG GGTGAACAAT TTTGTGGCTT TTTTCCCCCT ATAATTCTGA
CCCGCTCAGG CTTGAGGGTT TCTCCGGCCT CCGCTCACTG CGTGCACCTG
GCGCTGCCCT GCTTCCCCCA ACCTGTTGCA AGGCTTTAAT TCTTGCAACT
GGGACCTGCT CGCAGGCACC CCAGCCCTCC ACCTCTCTCT ACATTTTTGC
10 AAGTGTCTGG GGGAGGGCAC CTGCTCTACC TGCCAGAAAT TTTAAAACAA
AAACAAAAC AAAAAAATCT CCGGGGGCCC TCTTGGCCCC TTTATCCCTG
CACTCTCGT CTCTGCCCC ACCCGAGGT AAAGGGGGCG ACTAAGAGAA
GATGGTGTG CTCACGCGG TCTCCTGCT GCTGGCCGCC TATGCGGGG
CGGCCCAGAG CTTGGGCTCC TTCGTGCACT GCGAGCCCTG CGACGAGAAA
15 GCCCTCTCCA TGTGCCCCCC CAGCCCCCTG GGCTGCGAGC TGGTCAAGGA
GCCGGGCTGC GGCTGCTGCA TGACCTGCGC CTTGGCCGAG GGGCAGTCGT
GCGGCGTCTA CACCGAGCGC TGCGCCCAGG GGCTGCGCTG CCTCCCCCGG
CAGGACGAGG AGAAGCCGCT GCACGCCCTG CTGCACGGCC GCGGGGTTTG
CCTCAACGAA AAGAGCTACC GCGAGCAAGT CAAGATCGAG AGAGACTCCC
20 GTGAGCACGA GGAGCCCACC ACCTCTGAGA TGGCCGAGGA GACCTACTCC
CCCAAGATCT TCCGGCCCCA ACACACCCGC ATCTCCGAGC TGAAGGCTGA
AGCAGTGAAG AAGGACCGCA GAAAGAAGCT GACCCAGTCC AAGTTTGTCTG
GGGGAGCCGA GAACACTGCC CACCCCGGA TCATCTCTGC ACCTGAGATG
AGACAGGAGT CTGAGCAGGG CCCCTGCCGC AGACACATGG AGGCTTCCCT
25 GCAGGAGCTC AAAGCCAGCC CACGCATGGT GCCCCGTGCT GTGTACCTGC
CCAATTGTGA CCGCAAAGGA TTCTACAAGA GAAAGCAGTG CAAACCTTCC
CGTGGCCGCA AGCGTGGCAT CTGCTGGTGC GTGGACAAAGT ACGGGATGAA
GCTGCCAGGC ATGGAGTACG TTGACGGGA CTTTCAGTGC CACACCTTCG
ACAGCAGCAA CGTTGAGTGA TCGTCCCCC CCAAACCTTT CCCTCACCCC
30 CTCCACCCCC CAGCCCCGAC TCCAGCCAGC GCCTCCCTCC ACCCCAGGAC
GCCACTCATT TCATCTCATT TAAGGGAAAA ATATATATCT ATCTATTTGA
GGAAAAAAAA AAAAAAAAAA AA

```

(SEQ ID NO: 4)

35

These binding proteins can be made synthetically or cloned and produced for therapeutic purposes, while a cell line producing a desired monoclonal antibody can be maintained for relatively large-scale antibody production.

40

Cloning and general antibody methodologies are commonplace

in the art; such methodologies are disclosed within Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd ed. Cold Spring Harbor Laboratory Press 1989), the disclosure of which is hereby incorporated by reference as part of this disclosure.

Rather than using competitive methods, another aspect of the invention involves the use of a cholinergic special transporter to insert a gene which produces inactive receptors for one or more factor involved in promotion of neural sprouting. Such receptors would maintain high specific binding constants to their ligands, but the biological activity of the receptors would be abrogated; such receptors could be easily be generated and screened through the introduction of mutations in the nucleotide sequence encoding the protein, and assaying the mutants for binding strength and biological activity. Alternatively, the neurotrophic activities of factors produced by the neural endplate or nascent sprouts may be inhibited through intracellular targeting and delivery of a competitive inhibitor, ribozyme, transcriptional suppresser or other agent specifically able to block or attenuate such activities, as described above. Such intracellular targeting is disclosed in references such as, e.g., Dolly et al., International Patent Publication No. WO95/32738, previously incorporated by reference herein.

The second point at which intervention in the sprouting phenomenon may be made is during the stage of axonal outgrowth and arbor development. At this stage such

- outgrowth has already been initiated, but auxiliary factors appear to be necessary in order to maintain axonal growth. A number of factors are known to affect the rates of outgrowth, and may also effect the survivability of many
- 5 types of neurons. Such factors include ciliary neurotrophic factor (CNTF); neurotrophins, including NT-3, NT-4, and brain-derived neurotrophic factor (BDNF); and leukemia inhibitory factor (LIF). Not only are these factors important in establishing an initial rate of axonal
- 10 outgrowth, but they appear to either directly or indirectly stimulate their own production -- therefore blocking the initial outgrowth of sprouts may be essential in preventing further propagation through the continued expression of such factors.
- 15 The same methods as disclosed above can be used to prevent one or more of the agents listed above to manifest its activity. As would be expected by those of skill in the art in light of the present disclosure, such agents, destroy or bind to, and therefore attenuate the activity of these
- 20 factors or their block their ability to bind to their receptors, or inactive forms of their receptors, would be useful when used in conjunction with a clostridial neurotoxin to prevent or reduce the rate of sprouting of neurites from the poisoned or damaged endplate.
- 25 The third stage at which the sprouting phenomenon may be attenuated or inhibited concerns the binding of axons to the extracellular matrix. Axons are guided to cellular processes containing the appropriate neurotransmitter

receptors by binding to components of the extracellular matrix. Such binding involves a variety of cell-borne or matrix associated adhesion molecules. Tenascin-C is an extracellular matrix component derived from Schwann cells
5 that appears to bind neural processes. Ninjurin is a cell surface adhesion molecule that is up-regulated following peripheral nerve injury and thought to be involved in exonal guidance. See Araki, T., et al., *J. Biol. Chem.* 272:21373-21380 (1997), incorporated by reference herein. Similarly,
10 neural-cell adhesion molecule (N-CAM), is an adhesion molecule which is thought to be involved in binding of neural sprouts to the extracellular matrix. The same techniques indicated above may be employed to prevent the expression or inhibit the activity of these molecules when
15 used as part of a clostridial neurotoxin therapy.

Thus, in one aspect the present invention is drawn to a method for extending the effective period during which tissue treated with clostridial toxin is paralyzed, comprising: Contacting said tissue with a composition
20 comprising an agent able to prevent the neuroregenerative activity of a polypeptide selected from the group consisting of IGF-1, IGF-2, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule,
25 and neural agrin.

In one preferred embodiment, the agent comprises a polypeptide able to bind to IGF-1 and/or IGF-2 in a manner that prevents an IGF molecule from binding to or activating

a cell surface receptor involved in the initiation of neural sprouting. In a most preferred embodiment the polypeptide comprises at least a portion of a amino acid sequence selected from the group consisting of: IGFBP-4 (SEQ ID NO: 1) or IGFBP5 (SEQ ID NO: 2). Preferably said portion comprises at least 10 contiguous amino acids of said sequence; more preferably said portion comprises at least 20 contiguous nucleotides of said sequence. Most preferably, the portion comprises an amino acid sequence selected from the group consisting of the entire amino acid sequence of IGFBP-4 or IGFBP-5.

Treatment of cells with such a composition may be accomplished either before or simultaneously with treatment with clostridial toxin. Preferably, the clostridial toxin is a botulinum toxin. Even more preferably, the botulinum toxin comprises BoNT/A. In other embodiments, the clostridial toxin is TeNT. Agent which are able to bind to any of these factors in a manner that inhibits their neurotrophic activity, or which bind to the receptors for such factors, would, in light of the present application, be expected to function as agents for extending the effective period between treatments of tissue with a neurotoxin.

Another embodiment comprises a cholinergic specific transporter joined to a gene encoding a gene which produces an inactive receptor for one or more factor involved in promotion of neural sprouting when delivered to a neural cell in vivo. Preferably the receptors maintain high specific binding constants to said factor(s) and the

biological activity of the receptor is reduced or absent.

In a particularly preferred embodiment the transporter comprises some or all of a clostridial neurotoxin heavy chain, although other transporters such as the diphtheria toxin transporter may be effective in this regard as well.

In another embodiment the invention comprises a cholinergic specific transporter that is covalently or non-covalently joined to a nucleic acid which comprises a ribozyme or antisense nucleic acid able to specifically destroy the nucleic acids encoding neurotrophic agents or their receptors. Said joining may be made through methods including, but not limited to, covalent bonding or electrostatic forces.

In another embodiment, the present invention is drawn to the methods for stimulating the outgrowth of neural sprouts from damaged neural tissue. Such methods could be effective ways of increasing the rate at which reinnervation occurs after a neural injury. These methods comprise: Contacting said tissue with a composition comprising a polypeptide which comprises a neurotrophically active domain derived from an agent selected from a group consisting of IGF-1, IGF-2, ciliary neurotrophic factor, NT-3, NT-4, brain derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin. Such damage may be a result of neurotoxin poisoning or due to a traumatic event, including but not limited to nerve or spinal cord crush injuries, traumatic brain injuries, glaucoma-induced damage to the retina and/or

optic nerve, or surgical trauma or injury.

The following examples illustrate various embodiments of the present invention, and are not intended to limit the scope of the invention, which is solely defined by the claims which conclude this specification.

Example 1

Blepharospasm is a medical condition characterized by uncontrolled eyelid movement. In its early stages, the condition is characterized by excessive blinking or fluttering of the eyelids. The condition is generally a progressive one, in which excessive blinking is replaced in the later stages with spasms of eye closure that interfere with visual function. The spasms become more frequent and severe, and involve the preseptal, pretarsal, and orbicularis oculi muscle. The condition often results in functional blindness relatively quickly (in a matter of two to three years) after the symptoms are first encountered.

A patient suffering from moderate idiopathic blepharospasm is treated with injections of BoNT/A toxin preparation containing non-toxic proteins and hemagglutins in sterile saline. Alternatively, the same toxin preparation without hemagglutins may be used. The injections are generally in the volume of 100 μ l; and each injection contains 1.25 to 2.5 units of the toxin preparation. The injections are made into the pretarsal orbicularis oculi of the upper lid laterally and medially

and in the lower lid laterally and medially. Additionally,
2.5 unit injections (100 µl each) are made lateral to the
lateral canthus and into the brow medially. Total amount of
BoNT/A toxin injected is roughly 6.25 to 12.5 units per
5 eye. The Bo/A toxin is provided in a sterile, preservative-
free saline, and the same solution is used to dilute the
BoNT/A toxin if the master preparation of it is too
concentrated.

Following injection the treated muscles are
10 sufficiently paralyzed due to this treatment to alleviate
the major symptoms of blepharospasm. Some mild concomitant
weakness in the surrounding muscle tissue is observed; these
side effects are mild and tolerated well by the patient.
The effect of this treatment lasts approximately 8 weeks,
15 and must be repeated at the end of this time to maintain the
beneficial effects.

Example 2

A patient with blepharospasm is pre-treated with BoNT/A toxin as indicated in Example 1 with the following
5 difference. Prior to injection with BoNT/A toxin preparation, the patient is given a ten-fold excess of IGFBP-4, having an amino acid sequence of SEQ ID NO: 1. The binding protein preparation is dissolved in sterile, preservative-free saline. Each injection is in the same
10 area as the toxin injections that follow the pre-treatment; the volume of each injection is 100 µl. The BoNT/A toxin preparation is injected ten minutes after the injection of the IGFBT 4 injection.

The patient's therapeutic response to the BoNT/A toxin
15 is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended to 12 weeks or more, during which time no further injection need be made.

20

Example 3

A patient with blepharospasm is pre-treated with BoNT/A toxin as indicated in Example 1 with the following
difference. The BoNT/A toxin has been modified to have
25 joined thereto a nucleic acid comprising a ribozyme specifically targeted to enzymatically destroy neural agrin mRNA. No supplemental injections are made.

The patient's therapeutic response to the BoNT/A toxin is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended to 12 weeks or more, during which time no further injection
5 need be made.

Example 4

A patient with blepharospasm is pre-treated with BoNT/A
10 toxin as indicated in Example 1 with the following difference. The BoNT/A toxin has been modified to have joined thereto a nucleic acid encoding an inactive neurotrophin receptor which retains the ability to bind its target neurotrophin. No supplemental injections are made.

15 The patient's therapeutic response to the BoNT/A toxin is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended beyond that seen with BoNT/A alone, during which time no further injection need be made.

20 It will be understood that, while reference is made to BoNT/A in the Examples above, any other of the species of botulinum toxins (e.g., BoNT/B through G) could be substituted therefor, with appropriate adjustments possibly necessary due to differences in specific activity of the
25 toxin. Additionally, the light chain segment could be derived from any clostridial neurotoxin (or other neurotoxin) , with the heavy chain retaining the motor

neuron receptor binding and exo-vesicular transport
activities retained from the BoNT heavy chain.

These Examples are not intended to exhaust the
embodiments of the present invention, and the invention is
5 not to be seen as limited thereby. Further embodiments will
be disclosed within the claims that conclude this
specification.

CLAIMS

What is claimed is:

- 5 1) A method for extending the effective period during which tissue treated with a clostridial toxin is paralyzed comprising: contacting said tissue with a composition comprising an agent able to prevent the neuroregenerative activity of a polypeptide selected from the group consisting
10 of: IGF I, IGF II, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin.
- 15 2) The method of claim 1 wherein said contacting step occurs at the same time as said tissue is treated with said clostridial toxin.
- 3) The method of claim 1 wherein said contacting step
20 occurs prior to treatment of said tissue with said clostridial toxin.
- 4) The method of claim 1 wherein said clostridial toxin comprises BoNT.
- 25 5) The method of claim 1 wherein said clostridial comprises BoNT/A.

- 6) The method of claim 1 wherein said agent is selected from the group consisting of:
- a) an antibody able to selectively bind said polypeptide,
 - b) a competitive inhibitor of said polypeptide,
 - 5 c) a compound able to selectively prevent the expression of a gene encoding said polypeptide,
 - d) a binding protein other than an antibody, and
 - e) a ribozyme,
 - f) a nucleic acid encoding an inactive growth factor
- 10 receptor able to bind said growth factor.
- 7) The method of claim 6 wherein said agent is an antibody able to selectively bind said polypeptide.
- 15 8) The method of claim 6 wherein said agent is a competitive inhibitor of said polypeptide.
- 9) The method of claim 6 wherein said agent is a compound able to prevent the expression of a gene encoding said
- 20 polypeptide.
- 10) The method of claim 6 wherein said agent is a binding protein other than an antibody.
- 25 11) The method of claim 9 wherein said polypeptide is selected from the group consisting of IGF I and IGF II, and said binding protein is selected from the group consisting of IGF-BP4 and IGF-BP5.

- 12) A method for stimulating the outgrowth of neural sprouts from damaged neural tissue comprising: contacting said tissue with a composition comprising a polypeptide which comprises a neurotrophically active domain derived from an agent selected from the group consisting of IGF I, IGF II, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin.
- 13) The method of claim 11 wherein said agent comprises IGF I .
- 14) The method of claim 11 wherein said agent comprises IGF II.
- 15) The method of claim 11 wherein said agent comprises NT-3.
- 16) The method of claim 11 wherein said agent comprises ciliary neurotrophic factor.
- 17) The method of claim 11 wherein said agent comprises NT-3.
- 18) The method of claim 11 wherein said agent comprises NT-4.

19) The method of claim 11 wherein said agent comprises
brain-derived neurotrophic factor.

20) The method of claim 11 wherein said agent comprises
5 leukemia inhibitory factor.

21) The method of claim 11 wherein said agent comprises
tenascin-C.

10 22) The method of claim 11 wherein said agent comprises
ninjurin.

23) The method of claim 11 wherein said agent comprises
neural-cell adhesion molecule.

15

24) The method of claim 11 wherein said agent comprises
neural agrin.

SEQUENCE LISTING

<110> Dolly, Oliver J.
Aoki, Kei Roger
De Paiva, Anton

<120> Compositions and Methods for Modulating
Neural Sprouting

<130> 17259

<150> 60/083,472

<151> 1998-04-29

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 258

<212> PRT

<213> Homo sapiens

<400> 1

```

Met Leu Pro Leu Cys Leu Val Ala Ala Leu Leu Leu Ala Ala Gly Pro
 1           5           10           15
Gly Pro Ser Leu Gly Asp Glu Ala Ile His Cys Pro Pro Cys Ser Glu
          20           25           30
Glu Lys Leu Ala Arg Cys Arg Pro Pro Val Gly Cys Glu Glu Leu Val
          35           40           45
Arg Glu Pro Gly Cys Gly Cys Cys Ala Thr Cys Ala Leu Gly Leu Gly
          50           55           60
Met Pro Cys Gly Val Tyr Thr Pro Arg Cys Gly Ser Gly Leu Arg Cys
          65           70           75           80
Tyr Pro Pro Arg Gly Val Glu Lys Pro Leu His Thr Leu Met His Gly
          85           90           95
Gln Gly Val Cys Met Glu Leu Ala Glu Ile Glu Ala Ile Gln Glu Ser
          100          105          110
Leu Gln Pro Ser Asp Lys Asp Glu Gly Asp His Pro Asn Asn Ser Phe
          115          120          125
Ser Pro Cys Ser Ala His Asp Arg Arg Cys Leu Gln Lys His Phe Ala
          130          135          140
Lys Ile Arg Asp Arg Ser Thr Ser Gly Gly Lys Met Lys Val Asn Gly
          145          150          155          160
Ala Pro Arg Glu Asp Ala Arg Pro Val Pro Gln Gly Ser Cys Gln Ser
          165          170          175
Glu Leu His Arg Ala Leu Glu Arg Leu Ala Ala Ser Gln Ser Arg Thr
          180          185          190
His Glu Asp Leu Tyr Ile Ile Pro Ile Pro Asn Cys Asp Arg Asn Gly
          195          200          205
Asn Phe His Pro Lys Gln Cys His Pro Ala Leu Asp Gly Gln Arg Gly
          210          215          220
Lys Cys Trp Cys Val Asp Arg Lys Thr Gly Val Lys Leu Pro Gly Gly
          225          230          235          240
Leu Glu Pro Lys Gly Glu Leu Asp Cys His Gln Leu Ala Asp Ser Phe
          245          250          255

```

Arg Glu

<210> 2
 <211> 272
 <212> PRT
 <213> Homo Sapiens

<400> 2
 Met Val Leu Leu Thr Ala Val Leu Leu Leu Leu Ala Ala Tyr Ala Gly
 1 5 10 15
 Pro Ala Gln Ser Leu Gly Ser Phe Val His Cys Glu Pro Cys Asp Glu
 20 25 30
 Lys Ala Leu Ser Met Cys Pro Pro Ser Pro Leu Gly Cys Glu Leu Val
 35 40 45
 Lys Glu Pro Gly Cys Gly Cys Cys Met Thr Cys Ala Leu Ala Glu Gly
 50 55 60
 Gln Ser Cys Gly Val Tyr Thr Glu Arg Cys Ala Gln Gly Leu Arg Cys
 65 70 75 80
 Leu Pro Arg Gln Asp Glu Glu Lys Pro Leu His Ala Leu Leu His Gly
 85 90 95
 Arg Gly Val Cys Leu Asn Glu Lys Ser Tyr Arg Glu Gln Val Lys Ile
 100 105 110
 Glu Arg Asp Ser Arg Glu His Glu Glu Pro Thr Thr Ser Glu Met Ala
 115 120 125
 Glu Glu Thr Tyr Ser Pro Lys Ile Phe Arg Pro Lys His Thr Arg Ile
 130 135 140
 Ser Glu Leu Lys Ala Glu Ala Val Lys Lys Asp Arg Arg Lys Lys Leu
 145 150 155 160
 Thr Gln Ser Lys Phe Val Gly Gly Ala Glu Asn Thr Ala His Pro Arg
 165 170 175
 Ile Ile Ser Ala Pro Glu Met Arg Gln Glu Ser Glu Gln Gly Pro Cys
 180 185 190
 Arg Arg His Met Glu Ala Ser Leu Gln Glu Leu Lys Ala Ser Pro Arg
 195 200 205
 Met Val Pro Arg Ala Val Tyr Leu Pro Asn Cys Asp Arg Lys Gly Phe
 210 215 220
 Tyr Lys Arg Lys Gln Cys Lys Pro Ser Arg Gly Arg Lys Arg Gly Ile
 225 230 235 240
 Cys Trp Cys Val Asp Lys Tyr Gly Met Lys Leu Pro Gly Met Glu Tyr
 245 250 255
 Val Asp Gly Asp Phe Gln Cys His Thr Phe Asp Ser Ser Asn Val Glu
 260 265 270

<210> 3
 <211> 1955
 <212> DNA
 <213> Homo Sapiens

<400> 3
 gtgccctccg ccgctcgccc gcgcgcgcgc gctccccgcc tgcgcccagc gccccgcgcc
 60
 cgcgccccag tcctcgggcg gtcattgctgc cctctgcct cgtggccgcc ctgctgctgg
 120
 ccgcggggcc cgggcccagc ctgggcgacg aagccatcca ctgcccgcc tgctccgagg
 180

agaagctggc gcgctgccgc ccccccgtgg gctgcgagga gctgggtgcga gagccgggct
240
gcggctgttg cgccacttgc gccctgggct tggggatgcc ctgcggggtg tacaccccc
300
gttgcggctc gggcctgcgc tgctaccgc cccgaggggt ggagaagccc ctgcacacac
360
tgatgcacgg gcaaggcgtg tgcattggagc tggcggagat cgaggccatc caggaaagcc
420
tcgagccctc tgacaaggac gagggtgacc accccaacaa cagcttcagc ccctgtagcg
480
cccatgaccg caggtgcctg cagaagcact tcgccaaaat tcgagaccgg agcaccagt
540
ggggcaagat gaaggtcaat ggggcgcccc gggaggatgc cggcctgtg cccaggggct
600
cctgccagag cgagctgcac cgggcgctgg agcggctggc cgcttcacag agcgcaccc
660
acgaggacct ctacatcatc cccatcccca actgcgaccg caacggcaac ttccacccca
720
agcagtgtca cccagctctg gatgggcagc gtggcaagtg ctggtgtgtg gaccggaaga
780
cgggggtgaa gcttcggggg ggcctggagc caaaggggga gctggactgc caccagctgg
840
ctgacagctt tcgagagtga ggcctgccag caggccaggg actcagcgtc cctgtctact
900
cctgtgctct ggaggctgca gagctgacct agagtggagt ctgagtctga gtctgtctc
960
tgcctgcggc ccagaagttt ccctcaaagc cgcgtgtgca cgtgtgcgtg tgcgtgcgtg
1020
tgtgtgtgtt tgtgagcatg ggtgtgccct tggggtaagc cagagcctgg ggtgttctct
1080
ttggtgttac acagcccaag aggactgaga ctggcactta gccaagagg tctgagccct
1140
ggtgtgtttc cagatcgatc ctggattcac tcaactcactc attccttcac tcatccagcc
1200
acctaaaaac atttactgac catgtactac gtgccagctc tagttttcag ccttgggagg
1260
ttttattctg acttctctg attttggcat gtggagacac tcctataagg agagttcaag
1320
cctgtgggag tagaaaaatc tcattcccag agtcagagga gaagagacat gtaccttgac
1380
catcgtcctt cctctcaagc tagccagagg gtgggagcct aaggaagcgt ggggtagcag
1440
atggagtaat ggtcacgagg tccagacca ctcccaaagc tcagacttgc caggtccct
1500
ttctcttctt cccaggtcc ttcttttagg tctggttgtt gcaccatctg cttggttggc
1560
tggcagctga gagcctgct gtgggagagc gaagggggtc aaaggaagac ttgaagcaca
1620
gagggttagg gaggtggggc acatttctct gagcagtcag ggtgggaaga aagaatgcaa
1680
gagtgagctg aatgtgccta atggagaaga cccacgtgct aggggatgag gggcttctctg
1740
ggtcctgttc cctaccccat ttgtggtcac agccatgaag tcaccgggat gaacctatcc
1800
ttccagtggc tcgtccctg tagctctgcc tcctctcca tatctcctc ccctacacct
1860

ccctccccac acctccctac tcccctgggc atcttctggc ttgactggat ggaaggagac
1920
ttaggaacct accagttggc catgatgtct tttct
1955

<210> 4
<211> 1722
<212> DNA
<213> Homo sapiens

<400> 4
ggggaaaaga gctaggaaag agctgcaaag cagtgtgggc ttttccctt tttttgctcc
60
ttttcattac ccctcctccg ttttcacct tctccggact tcgcgtagaa cctgcgaatt
120
tcgaagagga ggtggcaaag tgggagaaaa gaggtgtag ggtttgggg tttttgttt
180
ttgtttttgt tttttaattt cttgatttca acattttctc ccacctctc ggctgcagcc
240
aacgcctctt acctgttctg cggcgccgcy caccgctggc agctgaggg tagaaagcgg
300
ggtgtatttt agattttaag caaaaatttt aaagataaat ccatttttct ccccccccc
360
caacgccatc tccactgcat ccgatctcat tatttcgggtg gttgcttggg ggtgaacaat
420
tttgtggctt ttttccctt ataattctga cccgctcagg cttgaggggt tctccggcct
480
ccgctcactg cgtgcacctg gcgctgccct gcttccccca acctgttgca aggctttaat
540
tcttgcaact gggacctgct cgcaggcacc ccagccctcc acctctctct acatttttgc
600
aagtgtctgg gggagggcac ctgctctacc tgccagaaat tttaaaaca aaacaaaaac
660
aaaaaaatct cggggggccc tcttggcccc tttatccctg cactctcgct ctctgcccc
720
accccgaggt aaagggggcg actaagagaa gatggtgttg ctcaccgagg tcctcctgct
780
gctggccgcc tatgcggggc cggcccagag cctgggctcc ttcgtgcact gcgagccctg
840
cgacgagaaa gccctctcca tgtgcccccc cagccccctg ggctgcgagc tggtaagga
900
gcggggctgc ggctgctgca tgacctgcgc cctggccgag gggcagtcgt gcggcgctca
960
caccgagcgc tgcgcccagg ggctgcgctg cctcccccg caggacgagg agaagccgct
1020
gcacgccctg ctgcacggcc gcggggtttg cctcaacgaa aagagctacc gcgagcaagt
1080
caagatcgag agagactccc gtgagcacga ggagcccacc acctctgaga tggccgagga
1140
gacctactcc cccaagatct tccggcccaa acacaccgc atctccgagc tgaaggctga
1200
agcagtgaag aaggaccgca gaaagaagct gaccagtcct aagtttgcg ggggagccga
1260
gaacactgcc ccccccgga tcctctctgc acctgagatg agacaggagt ctgagcaggg
1320
ccctgcccgc agacacatgg aggttccct gcaggagctc aaagccagcc cagcatgggt
1380

gccccgtgct gtgtacctgc ccaattgtga cgcgaaagga ttctacaaga gaaagcagtg
1440
caaacccttcc cgtggccgca agcgtggcat ctgctgggtgc gtggacaagt acgggatgaa
1500
gctgccaggc atggagtacg ttgacgggga ctttcagtgc cacaccttcg acagcagcaa
1560
cgttgagtga tgcgtcccc ccacaccttt cctcaccacc ctcaccacc cagccccgac
1620
tccagccagc gcctccctcc accccaggac gccactcatt tcatttcatt taagggaaaa
1680
atatatatct atctatttga ggaaaaaaaa aaaaaaaaaa aa
1722

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/08303

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/16 A61K38/18 A61K38/30 A61K38/17 A61K38/39
//(A61K38/16,39:395,38:17,31:70)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CARONI P. ET AL: "Signaling by insulin-like growth factors in paralyzed skeletal muscle: Rapid induction of IGF1 expression in muscle fibers and prevention of interstitial cell proliferation by IGF-BP5 and IGF-BP4" THE JOURNAL OF NEUROSCIENCE, vol. 14, no. 5, 1994, pages 3378-3388, XP002115584 cited in the application abstract page 3379, column 1, paragraph 2 page 3385, column 1 page 3386, column 2, paragraphs 2,3 --- -/--	1-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 September 1999

Date of mailing of the international search report

05/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/08303

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	CIFUENTES-DIAZ C.: "The peripheral nerve and the neuromuscular junction are affected in the tenascin-C deficient mouse" CELLULAR AND MOLECULAR BIOLOGY, vol. 44, no. 2, March 1998 (1998-03), pages 357-379, XP002115585 abstract pages 357-358 "introduction" ---	12,21
X	BOOTH C.M. ET AL: "An antibody to neural cell adhesion molecule impairs motor nerve terminal sprouting in a mouse muscle locally paralysed with botulinum toxin" NEUROSCIENCE, vol. 35, no. 1, 1990, pages 85-91, XP002115586 abstract page 85, column 1-2 ---	1-8,12, 23
X	GURNEY M.E. ET AL: "Induction of motor neuron sprouting in vivo by ciliary neurotrophic factor and basic fibroblast growth factor" THE JOURNAL OF NEUROSCIENCE, vol. 12, no. 8, 1992, pages 3241-3247, XP002115587 cited in the application the whole document ---	12,16
X	A. K. MCALLISTER ET AL: "Neurotrophin regulation of cortical dendritic growth requires activity" NEURON, vol. 17, 1996, pages 1057-1064, XP002115588 cited in the application the whole document ---	12,19
X	CARONI P. ET AL: "Role of muscle insulin-like growth factors in nerve sprouting: suppression of terminal sprouting in paralyzed muscle by IGF-binding protein 4" THE JOURNAL OF CELL BIOLOGY, vol. 125, no. 4, 1994, pages 839-902, XP002115589 cited in the application abstract page 897, paragraph 2 -page 898, paragraph 1 ---	1-8, 10-14

-/--

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 99/08303

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 721 215 A (AOKI KEI ROGER ET AL) 24 February 1998 (1998-02-24) column 2, line 20 - line 55 column 5: table 1 column 7; table 1 -----	1-9
T	PAIVA, A. ET AL: "Functional repair of motor endplates after botulinum neurotoxin type A poisoning: Biphasic switch of synaptic activity between nerve sprouts and their parent terminals" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, USA, vol. 96, March 1999 (1999-03), pages 3200-3205, XP002115590 cited in the application the whole document -----	1-24

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/08303

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-24 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11

Methods to extend the period during which a tissue treated with a Clostridial toxin is paralysed by contacting the tissue with the compositions mentioned in the said claims 1-11.

2. Claims: 12-24

Methods for stimulating the outgrowth of neural sprouts from damaged neural tissue, comprising contacting the tissue with a composition containing the agents specified in said claims 12-24.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/08303

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5721215 A	24-02-1998	AU 2341797 A	10-10-1997
		CA 2249196 A	25-09-1997
		EP 0889731 A	13-01-1999
		WO 9734624 A	25-09-1997
